

STA 360/602L: MODULE 6.2

BAYESIAN LINEAR REGRESSION (ILLUSTRATION)

DR. OLANREWAJU MICHAEL AKANDE

BAYESIAN LINEAR REGRESSION RECAP

- Sampling model:

$$Y \sim \mathcal{N}_n(\mathbf{X}\boldsymbol{\beta}, \sigma^2 \mathbf{I}_{n \times n}).$$

- Semi-conjugate prior for $\boldsymbol{\beta}$:

$$\pi(\boldsymbol{\beta}) = \mathcal{N}_p(\boldsymbol{\mu}_0, \boldsymbol{\Sigma}_0).$$

- Semi-conjugate prior for σ^2 :

$$\pi(\sigma^2) = \mathcal{IG}\left(\frac{\nu_0}{2}, \frac{\nu_0 \sigma_0^2}{2}\right)$$

FULL CONDITIONAL

$$\pi(\boldsymbol{\beta}|\mathbf{y}, \mathbf{X}, \sigma^2) = \mathcal{N}_p(\boldsymbol{\mu}_n, \boldsymbol{\Sigma}_n),$$

where

$$\boldsymbol{\Sigma}_n = \left[\boldsymbol{\Sigma}_0^{-1} + \frac{1}{\sigma^2} \mathbf{X}^T \mathbf{X} \right]^{-1}$$
$$\boldsymbol{\mu}_n = \boldsymbol{\Sigma}_n \left[\boldsymbol{\Sigma}_0^{-1} \boldsymbol{\mu}_0 + \frac{1}{\sigma^2} \mathbf{X}^T \mathbf{y} \right],$$

and

$$\pi(\sigma^2|\mathbf{y}, \mathbf{X}, \boldsymbol{\beta}) = \mathcal{IG} \left(\frac{\nu_n}{2}, \frac{\nu_n \sigma_n^2}{2} \right),$$

where

$$\nu_n = \nu_0 + n$$

$$\sigma_n^2 = \frac{1}{\nu_n} \left[\nu_0 \sigma_0^2 + (\mathbf{y} - \mathbf{X}\boldsymbol{\beta})^T (\mathbf{y} - \mathbf{X}\boldsymbol{\beta}) \right] = \frac{1}{\nu_n} \left[\nu_0 \sigma_0^2 + \text{SSR}(\boldsymbol{\beta}) \right].$$

SWIMMING DATA

- Back to the swimming example. The data is from Exercise 9.1 in Hoff.
- The data set we consider contains times (in seconds) of four high school swimmers swimming 50 yards.

```
Y <- read.table("http://www2.stat.duke.edu/~pdh10/FCBS/Exercises/swim.dat")
Y
```

```
##      V1  V2  V3  V4  V5  V6
## 1 23.1 23.2 22.9 22.9 22.8 22.7
## 2 23.2 23.1 23.4 23.5 23.5 23.4
## 3 22.7 22.6 22.8 22.8 22.9 22.8
## 4 23.7 23.6 23.7 23.5 23.5 23.4
```

- There are 6 times for each student, taken every two weeks. That is, each swimmer has six measurements at $t = 2, 4, 6, 8, 10, 12$ weeks.
- Each row corresponds to a swimmer and a higher column index indicates a later date.

SWIMMING DATA

- Given that we don't have enough data, we can explore hierarchical models. That way, we can borrow information across swimmers.
- For now, however, we will fit a separate linear regression model for each swimmer, with swimming time as the response and week as the explanatory variable (which we will mean center).
- For setting priors, we have one piece of information: times for this age group tend to be between 22 and 24 seconds.
- Based on that, we can set uninformative parameters for the prior on σ^2 and for the prior on β , we can set

$$\pi(\beta) = \mathcal{N}_2 \left(\mu_0 = \begin{pmatrix} 23 \\ 0 \end{pmatrix}, \Sigma_0 = \begin{pmatrix} 5 & 0 \\ 0 & 2 \end{pmatrix} \right).$$

- This centers the intercept at 23 (the middle of the given range) and the slope at 0 (so we are assuming no increase) but we choose the variance to be a bit large to err on the side of being less informative.

POSTERIOR COMPUTATION

```
#Create X matrix, transpose Y for easy computayion
Y <- t(Y)
n_swimmers <- ncol(Y)
n <- nrow(Y)
W <- seq(2,12,length.out=n)
X <- cbind(rep(1,n),(W-mean(W)))
p <- ncol(X)

#Hyperparameters for the priors
mu_0 <- matrix(c(23,0),ncol=1)
Sigma_0 <- matrix(c(5,0,0,2),nrow=2,ncol=2)
nu_0 <- 1
sigma_0_sq <- 1/10

#Initial values for Gibbs sampler
#No need to set initial value for sigma^2, we can simply sample it first
beta <- matrix(c(23,0),nrow=p,ncol=n_swimmers)
sigma_sq <- rep(1,n_swimmers)

#first set number of iterations and burn-in, then set seed
n_iter <- 10000; burn_in <- 0.3*n_iter
set.seed(1234)

#Set null matrices to save samples
BETA <- array(0,c(n_swimmers,n_iter,p))
SIGMA_SQ <- matrix(0,n_swimmers,n_iter)
```

POSTERIOR COMPUTATION

```
#Now, to the Gibbs sampler
#library(mvtnorm) for multivariate normal

#first set number of iterations and burn-in, then set seed
n_iter <- 10000; burn_in <- 0.3*n_iter
set.seed(1234)

for(s in 1:(n_iter+burn_in)){
  for(j in 1:n_swimmers){

    #update the sigma_sq
    nu_n <- nu_0 + n
    SSR <- t(Y[,j] - X%%beta[,j])%%(Y[,j] - X%%beta[,j])
    nu_n_sigma_n_sq <- nu_0*sigma_0_sq + SSR
    sigma_sq[j] <- 1/rgamma(1,(nu_n/2),(nu_n_sigma_n_sq/2))

    #update beta
    Sigma_n <- solve(solve(Sigma_0) + (t(X)%%X)/sigma_sq[j])
    mu_n <- Sigma_n %% (solve(Sigma_0)%%mu_0 + (t(X)%%Y[,j])/sigma_sq[j])
    beta[,j] <- rmvnorm(1,mu_n,Sigma_n)

    #save results only past burn-in
    if(s > burn_in){
      BETA[j,(s-burn_in),] <- beta[,j]
      SIGMA_SQ[j,(s-burn_in)] <- sigma_sq[j]
    }
  }
}
```

RESULTS

- Before looking at the posterior samples, what are the OLS estimates for all the parameters?

```
beta_ols <- matrix(0,nrow=p,ncol=n_swimmers)
for(j in 1:n_swimmers){
beta_ols[,j] <- solve(t(X)%*%X)%*%t(X)%*%Y[,j]
}
colnames(beta_ols) <- c("Swimmer 1","Swimmer 2","Swimmer 3","Swimmer 4")
rownames(beta_ols) <- c("beta_0","beta_1")
beta_ols
```

```
##           Swimmer 1   Swimmer 2 Swimmer 3   Swimmer 4
## beta_0  22.93333333  23.35000000  22.76667  23.56666667
## beta_1  -0.04571429   0.03285714   0.02000  -0.02857143
```

- Can you interpret the parameters?
- Any thoughts on who the coach should recommend based on this alone? Is this how we should be answering the question?

POSTERIOR INFERENCE

- Posterior means are almost identical to OLS estimates.

```
beta_postmean <- t(apply(BETA,c(1,3),mean))
colnames(beta_postmean) <- c("Swimmer 1","Swimmer 2","Swimmer 3","Swimmer 4")
rownames(beta_postmean) <- c("beta_0","beta_1")
beta_postmean
```

```
##           Swimmer 1  Swimmer 2  Swimmer 3  Swimmer 4
## beta_0 22.9339174 23.34963191 22.76617785 23.56614309
## beta_1 -0.0453998  0.03251415  0.01991469 -0.02854268
```

- How about credible intervals?

```
beta_postCI <- apply(BETA,c(1,3),function(x) quantile(x,probs=c(0.025,0.975)))
colnames(beta_postCI) <- c("Swimmer 1","Swimmer 2","Swimmer 3","Swimmer 4")
beta_postCI[,1]; beta_postCI[,2]
```

```
##           Swimmer 1 Swimmer 2 Swimmer 3 Swimmer 4
## 2.5%    22.76901  23.15949  22.60097  23.40619
## 97.5%   23.09937  23.53718  22.93082  23.73382
```

```
##           Swimmer 1  Swimmer 2  Swimmer 3  Swimmer 4
## 2.5%  -0.093131856 -0.02128792 -0.02960257 -0.07704344
## 97.5%  0.002288246  0.08956464  0.06789081  0.01940960
```

POSTERIOR INFERENCE

- Is there any evidence that the times matter?

```
beta_pr_great_0 <- t(apply(BETA,c(1,3),function(x) mean(x > 0)))
colnames(beta_pr_great_0) <- c("Swimmer 1","Swimmer 2","Swimmer 3","Swimmer 4")
rownames(beta_pr_great_0) <- c("beta_0","beta_1")
beta_pr_great_0
```

```
##           Swimmer 1 Swimmer 2 Swimmer 3 Swimmer 4
## beta_0      1.0000      1.0000      1.0000      1.0000
## beta_1      0.0287      0.9044      0.8335      0.0957
```

```
#or alternatively,
beta_pr_less_0 <- t(apply(BETA,c(1,3),function(x) mean(x < 0)))
colnames(beta_pr_less_0) <- c("Swimmer 1","Swimmer 2","Swimmer 3","Swimmer 4")
rownames(beta_pr_less_0) <- c("beta_0","beta_1")
beta_pr_less_0
```

```
##           Swimmer 1 Swimmer 2 Swimmer 3 Swimmer 4
## beta_0      0.0000      0.0000      0.0000      0.0000
## beta_1      0.9713      0.0956      0.1665      0.9043
```

POSTERIOR PREDICTIVE INFERENCE


- How about the posterior predictive distributions for a future time two weeks after the last recorded observation?

```
x_new <- matrix(c(1,(14-mean(W))),ncol=1)
post_pred <- matrix(0,nrow=n_iter,ncol=n_swimmers)
for(j in 1:n_swimmers){
  post_pred[,j] <- rnorm(n_iter,BETA[j,,]%*%x_new,sqrt(SIGMA_SQ[j,]))
}
colnames(post_pred) <- c("Swimmer 1","Swimmer 2","Swimmer 3","Swimmer 4")

plot(density(post_pred[, "Swimmer 1"]),col="red3",xlim=c(21.5,25),ylim=c(0,3.5),lwd=1.5,
      main="Predictive Distributions",xlab="swimming times")
legend("topleft",2,c("Swimmer1","Swimmer2","Swimmer3","Swimmer4"),col=c("red3","blue3",
lines(density(post_pred[, "Swimmer 2"]),col="blue3",lwd=1.5)
lines(density(post_pred[, "Swimmer 3"]),col="orange2",lwd=1.5)
lines(density(post_pred[, "Swimmer 4"]),lwd=1.5)
```

POSTERIOR PREDICTIVE INFERENCE

Predictive Distributions



POSTERIOR PREDICTIVE INFERENCE

- How else can we answer the question on who the coach should recommend for the swim meet in two weeks time? Few different ways.
- Let Y_j^* be the predicted swimming time for each swimmer j . We can do the following: using draws from the predictive distributions, compute the posterior probability that $P(Y_j^* = \min(Y_1^*, Y_2^*, Y_3^*, Y_4^*))$ for each swimmer j , and based on this make a recommendation to the coach.
- That is,

```
post_pred_min <- as.data.frame(apply(post_pred,1,function(x) which(x==min(x))))
colnames(post_pred_min) <- "Swimmers"
post_pred_min$Swimmers <- as.factor(post_pred_min$Swimmers)
levels(post_pred_min$Swimmers) <- c("Swimmer 1","Swimmer 2","Swimmer 3","Swimmer 4")
table(post_pred_min$Swimmers)/n_iter
```

```
##
## Swimmer 1 Swimmer 2 Swimmer 3 Swimmer 4
## 0.7790 0.0078 0.1994 0.0138
```

- Which swimmer would you recommend?

WHAT'S NEXT?

MOVE ON TO THE READINGS FOR THE NEXT MODULE!